

DUMPSBOSS.

HAAD Licensure Examination for Registered Nurses

HAAD HAAD-RN

Version Demo

Total Demo Questions: 10

Total Premium Questions: 150

Buy Premium PDF

<https://dumpsboss.co>

support@dumpsboss.co

support@dumpsboss.co
dumpsboss.co

QUESTION NO: 1

Whenever a child with thalassemia comes for blood transfusion, he is administered Desferoxamine (Desferal). The action of this drug is to:

- A. Inhibit the inflammatory process
- B. Enhance iron excretion
- C. Antagonize the effect of vitamin C
- D. Increase red blood cell production

ANSWER: B

QUESTION NO: 2

A patient is admitted to a hospital with acute renal failure. The patient wakes up complaining of abdominal pain. On assessment, the nurse observes edema to the patient's ankles and distended neck veins. The patient is dyspneic with a blood pressure of 200/96 mmHg. The proper nursing diagnosis for this patient is:

- A. Deficient fluid volume related to disease process
- B. Excess fluid volume related to decreased glomerular filtration rate
- C. Knowledge deficit related to proper medication regimen
- D. Acute pain related to renal edema

ANSWER: B

QUESTION NO: 3

The patient is receiving piperacillin (Pipril) I.V. Which of the following should the nurse consider when administering the medication to the patient?

- A. Assess for Stevens-Johnson Syndrome
- B. Be alert for the possibility of hypersensitivity
- C. Watch for signs of increasing severity of infection
- D. Instruct patient to increase the intake of protein-rich foods

ANSWER: B

QUESTION NO: 4

A 65-year-old patient is admitted with ischemic stroke. Which of the following would be initially assessed by the nurse to determine the patient's level of consciousness?

- A. Visual fields
- B. Deep tendon reflexes
- C. Auditory acuity
- D. Verbal response

ANSWER: D

QUESTION NO: 5

The nurse is preparing teaching plans for several patients. The nurse should recognize which of the following patients is at greatest risk for fluid and electrolyte imbalance?

- A. A 2-year-old patient who is receiving gastrostomy feedings
- B. A 20-year-old patient with a sigmoid colostomy
- C. A 40-year-old patient who is 3 days post-operative with an ileostomy
- D. A 60-year-old patient who is 8 hours post-renal arteriography

ANSWER: C

QUESTION NO: 6

A deficiency of which of the following vitamins can affect the absorption of calcium?

- A. Vitamin C
- B. Vitamin B6
- C. Vitamin D
- D. Vitamin B12

ANSWER: C

QUESTION NO: 7

Symptoms of alcohol withdrawal include:

- A. Euphoria, hyperactivity and insomnia

- B. Depression, suicidal ideation and hypersomnia
- C. Diaphoresis, nausea and vomiting and tremors
- D. Unsteady gait, nystagmus and profound disorientation

ANSWER: C

QUESTION NO: 8

Which of the following interventions should the nurse implement if a patient complains of cramps while irrigating the colostomy?

- A. Reduce the flow of solution
- B. Have the patient sit up in bed
- C. Remove the irrigation tube
- D. Insert the tube further into the colon

ANSWER: A

QUESTION NO: 9

A patient undergoes laminectomy. In the immediate post-operative period, the nurse should

- A. Monitor the patient's vital signs and log roll him to prone position
- B. Monitor the patient's vital signs and encourage him to ambulate
- C. Monitor the patient's vital signs and auscultate his bowel sounds
- D. Monitor the patient's vital signs, check sensation and motor power of the feet

ANSWER: D

QUESTION NO: 10

A registered nurse delegates care to a practical nurse. The person most responsible for the safe performance of the care is the:

- A. Head nurse who is in-charge of the unit
- B. The practical nurse assigned to provide the care
- C. The registered nurse who delegated the care to the practical nurse
- D. The nursing care coordinator who is the supervisor of the unit

ANSWER: C